

# Day 5: Disease Modeling on Networks with Node Dynamics

NME workshop  
July 8-13, 2013

Prof. Martina Morris  
Prof. Steven Goodreau  
Sam Jenness



Supported by the US National Institutes of Health

# Changing network size and composition

- As social networks change in size (say, for instance, as a village of  $n = 5,000$  nodes grows to  $n = 10,000$  nodes), which of the following do you think is generally preserved?
  - Number of edges?  $e$
  - Mean degree?  $2e/n$
  - Density?  $e/\binom{n}{2}$

# Changing network size and composition

- Applying the coefficients as is from a given *stergm* fit to a network of changing size will lead to preservation of density across time
- For one-mode networks: preserving mean degree instead requires a simple transformation of the edges coefficient in the formation model:

$$\theta_{new} = \theta_{old} - \ln(N_{old}) + \ln(N_{new})$$

- This is mathematically equivalent to partitioning the original edges term into an offset equal to  $\ln(N)$  and a residual, and then updating the offset as  $N$  changes.
- EpiNet handles this for you
- If you are going to code your own models outside EpiNet, you must handle this

# Changing network and composition

- As network composition changes, balancing will happen automatically – the explicit pairing of individuals requires this.
- Nevertheless – one does not always have straightforward control over this – e.g. with just an edges term in the model, two sexes will automatically “meet in the middle”
- Can change parameterizations to obtain different dynamic behavior – worth thinking through the behavior you expect, and what you see for your model
- Some theory to guide you can be found in Morris (1991), Koehly, Goodreau and Morris (2004), Krivitsky, Handcock and Morris (2011)

# Relational dissolution through death

- We fit our dynamic network using static data, with a process for dissolving relationships governed by a coefficient derived from relational duration
- All of this was done in a context that contained no information about death – another process that terminates relationships
- If we simply layer death on to our model (even with the size correction on the previous slide) our relationship durations and our number of relationships will start dropping down below the expected values we want
- Some aspects of this might be desired (e.g. if we could interview dead people we might find their past relationships to be shorter than those of the same birth cohort in our sample who are still alive), but others are likely not.

# Relational dissolution through death

- An approximate correction for this is:
  1. Calculate dissolution coefficients as before (without considering death)
  2. Estimate formation coefficients conditional on these dissolution coefficients.
  3. Calculate new dissolution coefficients that reflect the log-odds of a relationship sustaining conditional on both actors sustaining, which equals:

$$\text{logit} \left[ \frac{P(E_t) - P(N_t)}{P(\neg N_t)} \right]$$

where:

$P(E_t)$  = the overall prob. of a tie dissolving a time  $t$  from any cause =  $1/D$

$P(N_t)$  = the prob. of either incident node dying at time  $t$

# Relational dissolution through death

- The complicating factor is that  $P(N_t)$  may very well change over time as your disease prevalence changes
- But then again, it's probably OK for relationship lengths to shorten, and network density to decline, slightly as death increases
- Again, EpiNet handles this for you
- If you are going to code your own models outside EpiNet, must handle this
- Bigger point:

**DIAGNOSE THE HECK OUT  
OF YOUR SIMULATIONS!!!**

# Review of offsets and corrections

<p>When approximating the fit of a formation STERGM conditional on dissolution STERGM...</p>	<p>...subtract dissolution coefficients from corresponding formation ones</p>
<p>When network size N changes and you want to preserve mean degree...</p>	<p>...subtract the ln of the old N and add the ln of the new N to the edges coefficient in the formation model (or equivalently, use an edges offset and update it with ln of new N)</p>
<p>To adjust for deaths in simulating from a STERGM model estimated from a cross-sectional network and durations</p>	<p>...use</p> $\text{logit} \left[ 1 - \frac{P(E_t) - P(N_t)}{P(\neg N_t)} \right]$ <p>in calculating your dissolution coefficients</p>